

Literatur:

1. Schopper C., Moser D., Sabbas A., Spassova E., Goriwoda W., Lagogiannis G., Yerit K., Watzinger F., König F., Donath K., Ewers R.
The Fluorohydroxyapatite (FHA) Frios® Algipore® is a Suitable Biomaterial for the Reconstruction of Severely Atrophic Human Maxillae.
Clinical Oral Implants Research, Vol. 14 (2003)
2. Wanschitz F., Figl M., Wanger A., Ewers R.
Measurement of Graft Volume Changes after Sinus Floor Elevation with a Phycogene Hydroxylapatit.
Int J Oral Maxillofac Implants 2006; 21: 433-438
3. Rapone B, Inchingolo AD, Trasarti S, Ferrara E, Qorri E, Mancini A, Montemurro N, Scarano A, Inchingolo AM, Dipalma G, Inchingolo F.
Long-Term Outcomes of Implants Placed in Maxillary Sinus Floor Augmentation with Porous Fluorohydroxyapatite (Algipore® FRIOS®) in Comparison with Anorganic Bovine Bone (Bio-Oss®) and Platelet Rich Plasma (PRP): A Retrospective Study.
J. Clin. Med. 2022, 11, 2491: 1-14.
<https://doi.org/10.3390/jcm11092491>
4. Simunek A., Cierny M., Kopecka D., Kohout A., Bukac J., Vahalova D.
The sinus lift with phycogenic bone substitute – A histomorphometric study.
Clin Oral Impl Res, Vol.16, 3:342-348, 2005

5. Lee C.Y.S., Rohrer M.D., Prasad H.S., Stover J., Suzuki J.B.
Sinus grafting with a natural flurohydroxyapatite for immediate load.
J Oral Implantology 35(4), 2009: 164-175.
6. Poeschl P.W., Ziya-Ghazvini F., Schicho K., Buchta C., Moser D., Seemann R., Ewers R., Schopper C.
Application of Platelet-Rich Plasma for Enhanced Bone Regeneration in Grafted Sinus.
J Oral Maxillofac Surg 70, 2012: 657-664
7. Scarano A., Degidi M., Perrotti V. , Piattelli A., Iezzi G.
Sinus augmentation with phycogene hydroxyapatite: histological and histomorphometrical results after 6 months in humans. A case series.
Oral Maxillofac Surg 16(1) 2011 DOI 10.1007/s10006-011-0296-3, 1-5
8. Iezzi G., Degidi M., Piattelli A., Mangano C., Scarano A., Shibli J.A. , Perrotti V.
Comparative histological results of different biomaterials used in sinus augmentation procedures: a human study at 6 months. Clin Oral Impl Res 0, 2011: 1–8
9. Barbeck M., Najman S., Stojanovic S., Mitic Z., Zivkovic J. M., Choukroun J., Kovačević P., Sader R., Kirkpatrick CJ., Ghanaati S.
Addition of blood to a phycogenic bone substitute leads to increased in vivo vascularization.
Biomed Mater, 10(5): 2015, 055007. doi:10.1088/1748-6041/10/5/055007
10. Ewers R.
Maxilla Sinusgrafting with Marine Algae Derived Bone Forming Material: A Clinical report of long term results.

J Oral Maxillofac Surg, 63:1712-1723, 2005

11. Lindenmüller I.H., Lambrecht J.T.
Sinus floor elevation and implantation--a retrospective study. Schweiz Monatsschr Zahnmed. 2006;116(2):142-149
12. Khoury F, Keller P, Keeve PL.
Stability of Grafted Implant Placement Sites After Sinus Floor Elevation Using a Layering Technique: 10-Year Clinical and Radiographic Results.
Int J Oral Maxillofac Implants 32(5) 2017:1086-1096.
doi:10.11607/jomi.5832.
13. Sokolowsky A, Sokolowski A Schwarze U, Theisen K, Payer M, Lorenzoni M, Wegscheider W. Phycogenic bone substitutes for sinus floor augmentation:
Histomorphometric comparison of hydroxyapatite and biphasic calcium phosphate in a randomised clinical pilot study
Int J Oral Implantol; 13(4) 2020: 387–39